

3.3 Discussion

Potential Population Size. Potential population size is often estimated by dividing the area of xeric oak habitat by 10 ha. Dividing the area of xeric habitat by 10 ha resulted in a potential population size of 354 territories, which was similar to other estimates (e.g., Swain et al. 1995). Dividing our estimate of palmetto-oak by 10 ha suggested there was enough habitat for another 482 territories, which more than doubled the potential population size. Independent mapping studies by the Brevard County Natural Resources Department (2002) produced similar increases in estimated habitat. Therefore, the estimated amount of potential habitat has increased over time even though the actual amount of potential habitat has decreased because of rapid urbanization (Toland 2001). This occurred because our knowledge of species-specific habitat relationships has increased and because the application of land cover maps that are not specifically designed to map habitat of specialized species can result in many potential errors (Breininger et al. 1991a, 1995; Breininger and Oddy 2001). Landcover maps usually cannot distinguish secondary and tertiary territories from mesic flatwoods that is devoid of scrub oaks and is unsuitable. Our habitat mapping and independent mapping by Brevard County Natural Resources explicitly considered habitat features within the flatwoods matrix that were important to scrub species.

The relative distribution of potential territories among xeric oak and palmetto-oak changed when applying the potential territory model, which more realistically allocated territories along xeric oak scrub ridges. Within PRUs, the potential territory model suggested 535, 209, and 89 territories respectively for primary, secondary, and tertiary territories. Thus, the number of territories in xeric oak (primary) increased and the number allocated to palmetto-oak (secondary and tertiary) decreased. This occurred because we limited the total number of territories to the maximum amount possible based on the sum of total oak and palmetto-oak and because we preferentially allocated territories from primary to secondary to tertiary until the total number of potential territories equaled the total based on a sum of all xeric oak and palmetto-oak. The increased proportion of primary territories occurred because xeric oak scrub ridges were narrow relative to grid cells so that grid cells were not entirely comprised of xeric oak scrub allowing a greater allocation of primary territories than would be expected if primary territories would have needed to be entirely comprised of xeric oak scrub. Real primary territories overlapping scrub oak ridges along Florida's Atlantic Coast also are only partially comprised of xeric scrub oak patches and include many smaller poorly-drained ridges, palmetto-lyonia, marshes and other cover types (Breininger et al. 1991a, 1995, 1998).

When landscape features permitted, we subjectively extended our PRU boundaries a few hundred meters beyond xeric oak scrub and palmetto-oak to buffer the ridges to enhance dispersal (see Stith 1999), reduce edge effects (see Mumme et al. 2000, Bowman and Woolfenden 2001), enhance prescribed fire (see Breininger et al. 2002), and because actual Florida scrub-jay territory boundaries extend for hundreds of meters past the focal habitat patches (Breininger et al. 1995, 1998). Consequently, the PRUs might be able to support more tertiary territories than we estimated. A large number of tertiary territories could only be supported if there was a continuously large excess of potential breeders because tertiary territories are sinks (see section 5.0 and Breininger

and Oddy [2001]). This excess could only be sustained if habitat conditions were optimal and source territories exceeded sink territories (Breininger and Carter 2003). The proportion of a population that can be sustained in these areas will depend on the amount of sink habitat, the rate of population decline in sinks, and regional processes of productivity in sources and immigration into sinks (Dias 1996). This is further complicated by the cooperative breeding system and under what conditions jays choose to disperse into marginal habitat. Florida Scrub-Jays have propensities to avoid dispersing into some types of marginal habitat (Woolfenden and Fitzpatrick 1984, Fitzpatrick and Woolfenden 1986). The proportions of sinks that can be sustained by sources is best estimated using population models, because stage-based vital rates, environmental and demographic stochasticity, density dependence, dispersal propensities, catastrophes, and changing habitat conditions, make calculations too complicated using human intuition and simple math (Burgman et al. 1993, Akçakaya et al. 1999, Breininger et al. 2002). We kept our estimates of potential territories simple and conservative within PRUs in this report by using the maximum area of oak and palmetto-oak to provide a limit on the number of territories that could be supported within PRUs.

The proportion of primary, secondary, and tertiary territories that actually occur within landscapes can be estimated by the sample sizes for juvenile production in our study sites given that we attempted to quantify juvenile production in all territories within landscapes without preferentially studying jays in primary, secondary, or tertiary territories. These percentages were 63%, 31%, and 5% respectively for primary, secondary, or tertiary territories (Section 5, Table 9). The estimated percentages for the potential territory model were 64%, 25%, and 11% respectively for primary, secondary, or tertiary territories. These relative percentages were similar, although the proportion of territories occupied might have been greater if most habitat had not been of marginal habitat quality.

Regardless of these uncertainties, palmetto-oak allows the population to be much larger and better able to withstand environmental stochasticity (Pulliam 1988, Howe et al. 1991). This occurs because there is back-migration from suboptimal into optimal habitats (Breininger and Carter 2003, Breininger and Oddy unpublished manuscript). Palmetto-oak territories can also buffer oak scrub from human edge effects and they provide for greater connectivity.

Potential Conservation Areas. The population in 1992 was probably >400 pairs, which was the criteria for designating a core population (Stith et al. 1996). Therefore, the population could be designated a core population, making it one of the four largest populations within the species range. The potential habitat is sufficiently contiguous so that the three metapopulations could be considered as one based on Stith et al.'s (1996) criteria if habitat was restored and recolonized by Florida Scrub-Jays in areas that currently separate the metapopulations. Historical mapping indicates that the habitat was once contiguous (Duncan et al. unpublished data).

In Section 5.0 we document population exchange between the Central Brevard and South Brevard-Indian River-St Lucie metapopulations. Therefore, these could be considered the same metapopulation, which has a potential size that can exceed 400 pairs, although it might take time to reach such a recovery goal once enough habitat

was acquired and restored. Given rampant human population expansion from Palm Bay to Cocoa, it is also likely that the Viera territory cluster might become its own separate metapopulation that would be vulnerable to extinction. The small territory cluster at Wickam Park might supplement the Viera population because Wickam Park is a Brevard County park, although it lacks a specific conservation plan.

Historical aerial photographs indicate that most potential territories were open savannas and therefore should have been suitable Florida scrub-jay habitat (Breininger personal observation, Duncan and others unpublished data). Only about 1/2 of the potential habitat is protected if one assumed a recovery goal was to recover scrub-jay populations to the maximum extent possible without including suburban jays that seem to have poor long-term persistence probabilities (Stith et al. 1996, Breininger 1999, Bowman unpublished, Stith 1999). Almost half the potential territories within PRUs have not even been proposed by a conservation land acquisition organization for protection. Some of these include landscapes of scrub and flatwoods that are contiguous for >5.0 kilometers (e.g., north of Buck lake, south of Fellsmere Road).

Except for Central Brevard, these metapopulations have some of the greatest potential within the range of the Florida scrub-jay (Stith 1999). Much additional habitat protection is needed to reduce extinction risk (Root 1998, Breininger et al. 1999a, Stith 1999). Separated and greatly reduced metapopulations might be vulnerable to inbreeding and the reduction of genetic information necessary for adaptation and resistance to new threats. A new threat, for example, includes the introduced West Nile Virus.

Additional mitigation and habitat acquisition efforts need to focus on making subpopulations larger and enhancing the connectivity between metapopulations. These also should focus on reducing edge/area ratios, which do not influence Florida scrub-jay population viability (see Mumme et al. 1991, Bowman and Woolfenden 2001) but are also critical to consider for other species (Breininger et al. in press). Several potentially large PRUs need to be incorporated into land acquisition plans. Developing a plan for conservation is complicated by economics and the willingness of sellers (of land). Recovery planning must be opportunistic by setting targets for overall population size and arrangement that are flexible enough to be implemented on a regional scale where there is political resistance to growth management and land use planning specific to maintaining viable populations. The relevance of each PRU is described in Section 6.

Habitat Quality of Potential Territories.

Habitat quality of most potential territories was poor because tree densities had become too great or because shrubs had become too tall. These changes were becoming common by the 1960s as natural and prescribed fire (by ranchers) frequencies were reduced because of urbanization and fire suppression (Duncan et al. 1999, in press). Habitat quality was already poor by 1992 and probably had been poor for decades (Swain et al. 1995, Boyle 1996). Populations presumably have been declining for decades based on habitat quality and habitat-specific demographic data collected elsewhere (Woolfenden and Fitzpatrick 1991, Root 1998, Breininger et al. 1999a). Many conservation areas showed progress in restoring habitat, but most still

were suboptimal in 2002, because restoration is often a slow process (Schmalzer and Boyle 1998, Breininger and Carter 2003).

Habitat quality was generally worst in North Brevard because of a longer history of fire exclusion. Much scrub was still ranchland in Central and South Brevard during the 1970s and ranchers frequently used prescribed fire. Extensive fuels loads were reduced by 1998 wildfire in North Brevard, but these occurred in a region (Buck Lake and North) where Florida scrub-jay populations had declined to only a few pairs. Primary territories had a greater tendency to be tall with too many trees compared to secondary and tertiary territories because they do not burn as readily as palmetto-dominated habitats (Breininger et al. 2002). Primary territories should have had greatest propensities to be optimal, instead of short, if fire regimes were frequent (Breininger and Oddy unpublished manuscript).

Potential territories along roads tended to have the poor habitat quality not only because small habitat fragments have a lower propensity to burn (Duncan and Schmalzer unpublished) but also because edges themselves burn poorly (Breininger et al. 2002).

Population Size

Population declines were much greater than the rates of habitat loss, as expected because of poor habitat quality. The differences between actual population size and potential population size was probably greatest in North Brevard because habitat quality was the poorest and this probably occurred for the longest period because so many potential territories were tall and forested. Unless scrub has remained unburned for long periods, it takes an average of 20 years to become suboptimal and much longer to become forested (Schmalzer et al. 1994).

We do not know why population declines did not appear in Central Brevard between 1992 and 2002. One possibility was that 1992 populations were underestimated and another possibility is that habitat quality actually improved between 1992 and 1999. Habitat quality was generally poor in 1992 (Swain et al. 1995, Boyle 1998) but wildfires and habitat management at the Viera mitigation site caused improvements. Site-specific descriptions of population size changes are provided elsewhere (Brevard County Natural Resources Department 2002).

4.0 Demography and Dispersal of Florida Scrub-Jays on the Mainland of Central Florida's Atlantic Coast.

The focus was to classify actual territories based on their habitat characteristics and quantify their demography. The quantification of habitat-specific demography is critical for understanding how habitat arrangements influence population dynamics and for developing population management strategies (Pulliam et al. 1992). We focus on the territory scale because these are the most basic landscape unit for quantifying habitat-specific population processes (Breininger and Carter 2003). We compare habitat potential of primary, secondary, and tertiary territories described in Chapter 3 to evaluate the potential contributions of palmetto-oak to Florida scrub-jay populations. We evaluate whether the edges of conservation reserves are likely to be population

sinks, which would suggest extirpation of all populations, except in the largest reserves. We test rapid habitat assessment procedures that might explain the population decline and be used to determine prescribed fire management needs. We do not use our more detailed habitat suitability index model (Duncan et al. 1995, Breininger et al. 1998) because it is cumbersome for managers to implement, especially given the uncertainties (Burgman et al. 2001). We also quantify dispersal patterns and exchanges among territory clusters that we observed.

4.1 Methods

Study Sites. On the mainland, colorbanding began during 1997 in South Brevard because most Florida Scrub-Jays and protected areas occurred there. We expanded study sites as areas were acquired for conservation allowing us to obtain access for the following investigations. A Florida Fire Science Team was formed to address questions that remained following the large 1998 wildfires. This team funded our initial expansion into North Brevard (Breininger et al. 2001). In 2000, we expanded into Central Brevard because of rapid habitat destruction and concerns by the U.S. Fish and Wildlife Service and recovery team biologists. The study ended prematurely after July 2002 because of funding. Figure 17 and Table 7 summarize the locations of territory mapping, demography, and dispersal studies involving color-banded Florida Scrub-Jays. We did not perform demography studies at Hollywood and Pirates territory clusters, although territory boundaries were provided. To quantify population size and dispersal into study tracts, we banded Florida Scrub-Jays at many sites where we did not perform territory mapping or demography studies (e.g., Port St John, East Micco, and Wabasso). Previous sections described that surveys for dispersed jays were conducted throughout 85% of the metapopulation.

Table 7. Number of Florida scrub-jay territories investigated on the mainland of Brevard County and Indian River County.

Territory Cluster	1997	1998	1999	2000	2001	2002
Buck Lake	0	0	3	2	2	2
Seminole Ranch	0	0	0	11	8	7
Tico-Grissom	0	0	0	16	20	17
Viera	0	0	0	1	18	14
Wickam	0	0	0	10	17	17
Malabar-Jordan-Valkaria	35	25	28	33	33	43
Stewart	0	0	0	0	4	4
Palm Bay	26	17	13	18	20	19
West Micco	13	11	9	9	10	11
Sebastian	3	3	3	16	14	14
Fellsmere	5	5	6	4	31	32
Totals	82	61	62	120	177	180

Capture, colorbanding, and census. For each new study area, we repeatedly surveyed the site 2-7 days each week to quantify the number of Florida scrub-jay territories and individuals occupying the site. During this time, we exposed Scrub-Jays to raw peanut bits to prepare them for capture. Once jays became familiar with peanuts, we baited Potter traps but kept the trap doors wired open. Once many jays entered traps, we began the capture process. In addition to Potter traps, we used drop traps to capture jays unwilling to enter Potter traps. Mist nets were also used to capture jays unwilling to enter either trap. Captured jays were banded with a unique combination of one silver and 2-3 colorbands. Once captured, covert feathers were examined to distinguish individuals less than one-year-old from older jays. We attempted to capture every jay providing we had access to its territory. Unbanded jays remained at most sites because they were difficult to capture or because we could not obtain permanent access. Conservation often involved partial acquisition of potential habitat within landscapes and this influenced our access. Access improved during the study as more land was acquired.

Throughout the study we color-banded juveniles and immigrants. We had difficulties capturing many individuals on several sites so that some demographic analyses (e.g., survival) involved samples of the population at those sites and not complete demographic analyses of every territory. Most Florida Scrub-Jays were color-banded at a site before we began quantifying survival at the site. Breeders were identified based on dominance behaviors, intensity of territory defense, and nesting activities (Woolfenden and Fitzpatrick 1984, 1991, 1996; Breininger et al. 1996a; Breininger 1999). Our efforts were to focus on banding as many jays in as many territories as possible, rather than conducting the intensive systematic nest searches sometimes performed. Juveniles were tallied among all territories in July. We sometimes quantified juvenile production before most individuals were color-banded because juveniles in July were conspicuous and nearly always could be found with their families.

Territory mapping. Territory mapping was conducted from April through May. Normally, boundaries were delineated to within a few meters by initiating disputes between families (Woolfenden and Fitzpatrick 1984, Breininger et al. 1995). All published studies have involved areas where all scrub was rigorously contested and occupied by jays (Woolfenden and Fitzpatrick 1984, Breininger et al. 1995, 1998). In this study, territory disputes did not occur in some large expanses of mesic flatwoods, extensively burned areas, forests, agricultural areas, or urban areas that were adjacent to areas occupied by Florida Scrub-Jays. Areas of unoccupied suitable habitat often occurred adjacent to occupied territories. In these cases, we mapped boundaries based on areas where jays appeared unwilling to enter. Territory boundaries for every year were digitized as ARC/INFO polygon files. The primary purpose of these boundaries was to characterize habitat quality at the territory scale. We did not focus our analyses on territory size because Florida Scrub-Jays used much larger areas than they need because population sizes were far below carrying capacity in most areas.

Territory quality. Oak cover and proximity to human edges represented habitat potential; height arrangements represented current habitat suitability and were influenced by fire history. Oak cover in territories was classified as primary if the territory intersected ≥ 0.4 ha of oak scrub on well-drained soils (Breininger and Oddy 2001). We used U.S. Department of Agriculture soils maps (Huckle et al. 1974, Wettstein et al. 1984) to define well-drained soils. Territories were designated secondary if they did not intersect well-drained oak scrub but intersected scrub oak polygons having $> 50\%$ scrub oak cover but ≥ 0.4 ha. Territories were designated tertiary if they lacked any oak scrub polygons > 0.4 ha in size that had $> 50\%$ oak cover. We classified territories into three categories based on their context to human-dominated landscapes. "Core" territories were not adjacent to human housing or hard surface roads. "Edge" territories were within potential reserves and bordered human landscapes, but were not adjacent to hard surface roads. "Suburban territories" occurred within suburban areas or were adjacent to hard surface roads. Height was classified using criteria in Table 3 derived from long-term studies at KSC (Breininger and Carter 2003).

Figure 17. Locations of Florida scrub-jay territories that were mapped in 2001 or 2002. Florida Scrub-Jays were also colorbanded in many other locations.



Analyses. Individual study years were from 1 April to 31 March. The demographic study funded by the U. S. Fish and Wildlife Service ended in July 2002 because of funding cuts. We did not separate the 1997 epidemic year from other years because we no longer assumed epidemics were rare annual events (see Woolfenden and Fitzpatrick 1984). West Nile Virus is a new potential disease that could be common and widespread, as confirmed in sentinel chickens with Brevard County and Indian River County during the summer of 2002. Related studies in 2002 suggest that survival of Brevard and Indian River county Scrub-Jays was greatly reduced by this new disease.

Continuities of pair-bonds between successive breeding seasons were compiled for all years for pairs where both breeders were colorbanded. Four categories were used to describe changes in pair-bonds between breeding seasons: one breeder paired with another Florida Scrub-Jay after the death of its spouse in the same territory; one breeder became a nonbreeder after the death of its spouse; mortality of both breeders; and divorce (Breininger et al. 1996a, Breininger 1999).

We calculated the mean distance between natal and breeding destination territories for males and females (see Woolfenden and Fitzpatrick 1984). We also developed a dispersal curve that was based on the number of territories that occurred between the natal territory and the territory where jays first bred. This was produced by establishing a straight line between natal and breeding destination territories and then counting the actual territories that intersected the line. We compiled all data on Florida Scrub-Jays having a known history since hatching to quantify the frequency that one-year-old jays delayed breeding.

We defined juveniles as young Florida Scrub-Jays present in July when most of them approached nutritional independence (Woolfenden and Fitzpatrick 1984). For each year, the number of juveniles and yearlings produced per breeding pair was calculated in every territory.

Annual survival for helpers and breeders was calculated as the number of jays that survived during a study year divided by the number alive at the beginning of the study year (Woolfenden and Fitzpatrick 1984, 1996; Breininger et al. 1995, 1996a, 1998a). We calculated birds as having survived if they were seen anytime after the annual survival period ending date. Survival analyses were performed only on colorbanded individuals. Bowman (unpublished report) compared these approaches with mark-recapture models and determined that breeder survival analyses produced identical results but that the above approaches produced slight, but insignificantly lower, nonbreeder survival rates. We didn't perform mark recapture analyses because our mean monthly detection probabilities exceeded 0.9 and so that our apparent survival estimates approached true survival estimates, excluding permanent and undetected emigration. To investigate potential errors we tallied all birds that were known to be alive but that were not observed for an entire year. We also tallied all jays known to have emigrated but that would not have been known to survive had we not been surveying most of the population on a regular basis. Extensive peripheral surveys allowed us to observe most successful offsite dispersals because we surveyed >85% of the population. We will investigate these small errors more extensively using multi-state models (Williams et al. 2001) and other approaches (e.g., Koenig et al. 2000) as we prepare results for journal publication.

We calculated demographic performance for every year in every territory within our study area by subtracting the number of breeders that died from the number of yearlings that were produced (Breininger et al. 1995, 1998). Demographic performance was calculated only for pairs where both breeders were colorbanded. We assumed that our samples approximated independent observations because there was more than normal annual breeder turnover, because individual territories often varied among treatment categories (e.g., breeder experience, presence/absence of helpers), because there was much annual variation in spatial distributions of habitat suitability, and because there was annual variation in environmental factors (e.g., rainfall) and population density. Furthermore, study site expansions allowed us to increasingly sample a larger portion of the actual population so that new pairs were continuously being added into the sample and we studied a large proportion of the population.

We used one-sample Kolmogorov-Smirnoff tests to determine whether data differed from a normal distribution. If they differed significantly ($p < 0.05$), we used nonparametric Mann Whitney U exact tests for two treatment tests and Kruskal-Wallis exact tests for multiple treatment tests (SPSS 1999). We also used ANOVAs (SPSS 1999) to test whether means varied among treatments (habitat types, presence of helpers). When assumptions regarding normality and the homogeneity of variance were violated, we still focused our presentation on the parametric ANOVAs, providing the conclusions regarding the results of statistical significance were the same ($p < 0.05$) with the nonparametric tests. We used parametric analyses of variances because they tend to be robust to deviations from normality and homogeneity of variances and we wanted to perform pair-wise comparisons. Providing the means were significantly different, pair-wise comparisons were performed using Tukey's tests when variances were homogeneous. We used the Games-Howell tests when variances were not equal (SPSS 1999). Differences in breeder survival among treatments categories were tested using likelihood ratio exact tests.

More complex analyses will be performed on a set of alternative general linear models to test for interactions among variables as the data is prepared for scientific journal publication. We chose to avoid such analyses in this report in order to avoid the negative consequences of data dredging and because preliminary analyses revealed that limited sample sizes for optimal territories greatly limited the management utility of multivariate analyses. Sample sizes for optimal territories were only beginning to become sufficient for investigating multivariate relationships prior to termination of the study. A slow accumulation of optimal territories across a range of other habitat factors occurred because there were few optimal territories in public ownership when the studies began. The number of optimal territories increased slowly because there was a lag between the time habitat became optimal after land acquisition because of vegetation response to fire and the logistical difficulties associated with restoration (Breininger and Carter 2003).

4.2 Results

We color-banded 800 Florida Scrub-Jays and studied demography of 180 Florida scrub-jay territories during 2002, which represented approximately 64% of the combined North Brevard, Central Brevard, and South Brevard-Indian River-St. Lucie metapopulations. We present 2 examples to demonstrate the dynamics of habitat occupancy and territory shifts. The first example shows that the amount of habitat occupied can change greatly across years and that Florida Scrub-Jays were not restricted to xeric oak scrub (Figure 18). In this example, all palmetto-oak territories were almost always occupied but xeric oak territories were often not occupied because they did not burn enough or they burned too extensively. The second example shows an example of variation between years (Figure 19). None of the examples were unusual for the data set.

Basic Demography. Of pair bonds where both breeders were color-banded, 61% stayed together between successive years, 23% involved the death of one breeder and the survivor finding a replacement breeder, 11% involved the death of both breeders, 4% involved the death of one breeder with the survivor becoming a non-breeder, and 1% resulted in divorce. The respective probabilities that individual breeders bred, helped, or floated during the next year were 0.97, 0.02, and 0.01 for breeders that survived the year ($n = 754$).

Mean family size during the study was 2.77 adults ($n = 682$). The respective probabilities that helpers bred, helped, or floated during the next year were 0.36, 0.62, and 0.02 for helpers that survived to the following year ($n = 254$). Sixty-five percent of female one-year-olds ($n = 101$) helped during the first nesting season after hatching. Seventy-nine percent of male one-year-olds ($n = 136$) helped during the first nesting season after hatching. Sex-specific differences in the frequency of helping by one-year-olds were significant (log likelihood ratio value = .023; $P = 5.185$).

A total of 312 color-banded jays permanently disappeared. There were 7 Florida Scrub-Jays that disappeared for >1 year and were later re-sighted. These included 3 breeders, 3 juveniles, and 1 helper. Only 2 of these did not move to another territory cluster. Only 6 breeders were observed to move from the cluster where they were observed to have first bred. Only 3 one-year olds moved to a territory cluster that was different than their natal cluster. Thirteen helpers moved to another territory cluster.

There was much annual variation in demography (Table 8). Female breeder survival was 0.75 ($n = 285$) and male breeder survival was 0.76 ($n = 331$); these differences were not significant (log likelihood ratio value = .151; $P = 0.768$). Female helper survival was 0.76 ($n = 106$) and male helper survival was 0.73 (103) these differences were not significant (log likelihood ratio value = .361; $P = 0.548$).

Juvenile production was 0.94 ($n = 190$) for experienced breeders and 0.52 ($n = 56$) for novice breeders; these differences were significant ($F = 4.527$; $P = 0.034$). Yearling production was 0.58 ($n = 111$) for experienced breeders and 0.40 ($n = 40$) for novice breeders; the P value for these differences was 0.305 ($F = 1.061$). Survival of experienced breeders was 0.81 ($n = 303$), and survival of novice breeders was 0.73 ($n = 86$); the P value for these differences was 0.134 (log likelihood ratio value = 2.248).

Demographic performance was 0.23 ($n = 108$) for experienced breeders and -0.26 ($n = 38$) for novice breeders; these differences were significant ($F = 4.659$; $p = 0.033$).

Juvenile production was 0.94 ($n = 335$) for breeders with helpers and 0.66 ($n = 347$) for pairs without helpers; these differences were significant ($F = 8.593$; $P = 0.004$). Yearling production was 0.56 ($n = 240$) for breeders with helpers and 0.42 ($n = 255$) for pairs without helpers; the P value for these differences was 0.094 ($F = 2.807$). Survival was 0.80 for breeders with helpers and 0.72 for breeders without helpers; these differences were significant ($F = 6.020$; $P = 0.014$). Demographic performance was 0.214 ($n = 135$) for pair with helpers and -0.13 ($n = 127$) for pairs without helpers; these differences were significant ($F = 5.342$; $P = 0.022$).

Demography and Habitat Arrangements. A large number of territories occurred in palmetto-oak even and most territories not located along xeric oak scrub (primary ridges) occurred in secondary territories (Table 9). Secondary territories did not have lower demographic success than primary territories; tertiary territories were population sinks.

Table 8. Annual variation in demography and comparison with other published studies

Location	Year	Juveniles /pair	Yearlings /pair	Demographic performance	Breeder Survival	Helper Survival
Brevard Mainland	1997	0.83	0.39	-0.82	0.51	0.77
Brevard Mainland	1998	0.65	0.36	-0.09	0.73	0.78
Brevard Mainland	1999	1.75	0.80	0.35	0.72	0.75
Brevard Mainland	2000	0.74	0.32	-0.05	0.77	0.68
Brevard Mainland	2001	0.82	0.59	0.20	0.83	0.77
Brevard Mainland	2002	0.52	N/a	N/a	N/a	N/A
Mean of Annual Means		0.89	0.49	-0.08	0.71	0.75
South Brevard Beaches ^a	1993- 1997	0.41	0.22	Not reported	0.73	0.48
Happy Creek at KSC ^b	1989- 1993	0.47	0.32	Not reported	0.80	0.73
Tel 4 at KSC ^b	1988- 1993	0.96	0.62	Not reported	0.76	0.72
Optimal Scrub at Archbold ^c	1969- 1986	1.23	0.68	Not reported	0.81	0.74
Tall Scrub at Archbold ^d	1969- 1986	0.80	0.36	Not reported	0.72	Not reported

^a Breininger (1999).

^b Breininger et al. 1996a

^c Woolfenden and Fitzpatrick 1984

^d Woolfenden and Fitzpatrick 1991

Figure 18. Comparison of Occupied Territories at Valkaria.






Figure 19. Comparison of Occupied Territories at Palm Bay.



- 2000 Territory Boundaries
- 1999 Territory Boundaries
- Scrub
- Xeric Oak
- Palmetto-Oak
- Palmetto



Caption 9. Juvenile Florida Scrub-Jays showing brown head and silver tips on coverts.



Table 9. Demography among territories having different scrub oak cover.

	Type of scrub (mean \pm standard error; sample sizes in parentheses)			Statistics	
	Primary ^a	Secondary ^b	Tertiary ^c	ANOVA (F)	P
Juvenile Production (pairs)	0.85 \pm 0.06 (432)	0.75 \pm 0.08 (213)	0.43 \pm 0.12 (37)	2.235	0.108
Yearling Production (pairs)	0.51 \pm 0.06 (309)	0.47 \pm 0.81 (161)	0.32 \pm 0.12 (25)	0.562	0.571
Demographic Performance (pairs)	-0.02 \pm 0.09 ^{e, f} (162)	0.26 \pm 0.12 ^e (86)	-0.50 \pm 0.27 ^f (14)	3.070	0.048
Breeder Survival	0.73 \pm 0.02 (381)	0.84 \pm 0.03 (207)	0.59 \pm 0.09 (27)	12.635 ^d	0.002

^a = xeric oak scrub.

^b = palmetto-oak with patches of oak scrub \geq 0.4 ha and no xeric oak scrub.

^c = palmetto-oak with no patches of oak scrub >0.4 ha and no xeric oak scrub.

^d = Log likelihood test value; ANOVA not performed on breeder survival.

^{e, f} = Post Hoc analyses suggested these were not significantly different $p > 0.05$.

Proximity of territories to human landscapes was not a good predictor of demographic success when other habitat factors were not considered (Table 10). The arrangement of shrub height within territories was the most important predictor of demographic success (Table 11). Sample sizes were low for short territories during all years. Sample sizes were also low for optimal territories, except during 2001 and 2002. Only optimal scrub produced an excess of potential breeders. Although sample sizes were too low to draw conclusions, reserve edges did not appear to be population sinks if habitat conditions approached optimal (Table 12). We just began accumulating samples to investigate the influence of human landscape context in territories with optimal height arrangements when funding expired.

Table 10. Demography among territories having different attributes of human landcover.

	Type of scrub (mean \pm standard error; sample sizes in parentheses)			Statistics	
	Core ^a	Reserve Edges ^b	Suburbs ^c	ANOVA (<i>F</i>)	<i>P</i>
Juvenile Production	0.88 \pm 0.08 (277)	0.90 \pm 0.13 (108)	0.69 \pm 0.60 (297)	2.195	0.112
Yearling Production	0.58 \pm 0.07 (187)	0.59 \pm 0.10 (80)	0.38 \pm 0.06 (228)	2.984	0.052
Demographic Performance	-0.06 \pm 0.13 (104)	0.27 \pm 0.18 (51)	-0.08 \pm 0.10 (107)	1.553	0.214
Breeder Survival	0.74 \pm 0.03 (232)	0.79 \pm 0.04 (112)	0.76 \pm 0.03 (271)	1.439 ^d	0.487

^a = not adjacent to human-dominated landscapes.

^b = in potential reserves but adjacent to human-dominated landscapes but not hard surface roads.

^c = surrounded by human landcover types or bordering hard surface roads.

^d = log likelihood test value; ANOVA not performed on breeder survival.

Table 11. Demography among territories having different fire history (shrub height arrangements).

	Type of scrub (mean \pm standard error; sample sizes in parentheses)				Statistics	
	All Short ^a	Optimal ^b	Tall Mix ^c	All Tall ^d	ANOVA (F)	P
Juvenile Production	0.70 \pm 0.40 (10) ^{f,g}	1.35 \pm 0.14 (121) ^f	0.70 \pm 0.06 (367) ^{f,g}	0.64 \pm 0.08 (184) ^g	10.173	0.000
Yearling Production	0.22 \pm 0.22 (9) ^f	1.17 \pm 0.18 (64)	0.46 \pm 0.05 (274) ^f	0.26 \pm 0.05 (148) ^f	16.296	0.000
Demographic Performance	-0.38 \pm 0.26 (8) ^f	0.82 \pm 0.22 (38)	0.03 \pm 0.11 (144) ^f	-0.30 \pm 0.113 (72) ^f	8.172	0.000
Breeder Survival	0.69 \pm 0.12 (16)	0.88 \pm 0.03 (88)	0.76 \pm 0.02 (334)	0.71 \pm 0.03 (177)	9.949 ^e	0.019

^a = entire territory was < 120 cm tall except for scattered trees and shrubs.

^b = entire territory < 170 cm tall except for scattered trees and shrubs and at least 0.13 ha oak at 120-170 cm tall among short scrub .

^c = territory included at least 0.4 ha of tall shrubs (170 cm) among shorter shrubs.

^d = entire territory was > 170 cm tall except for some shrubs.

^e = log likelihood test value; ANOVA not performed on breeder survival.

^{f,g} = Post Hoc analyses reveal that these are not significantly different $p < 0.05$.

Caption 10. Example of another secondary ridge.



Table 12. Demography among primary and secondary territories having optimal height but different attributes of human landcover.

	Type of scrub (mean \pm standard error; sample sizes in parentheses)			Statistics	
	Core ^a	Reserve Edges ^b	Suburbs ^c	ANOVA (F)	P
Juvenile Production	1.40 \pm 0.15 (81)	1.54 \pm 0.53 (13)	1.11 \pm 0.34 (27)	0.458	0.634
Yearling Production	1.25 \pm 0.18 (40)	1.25 \pm 0.41 (8)	0.94 \pm 0.52 (16)	0.279	0.757
Demographic Performance	0.83 \pm 0.27 (24)	1.43 \pm 0.43 (7)	0.14 \pm 0.51 (7)	1.696	0.198
Breeder Survival	0.86 \pm 0.05 (57)	1.00 \pm 0.00 (12)	0.84 \pm 0.08 (19)	3.499 ^d	0.174

^a = not adjacent to human-dominated landscapes.

^b = in potential reserves but adjacent to human-dominated landscapes but not hard surface roads.

^c = surrounded by human landcover types or bordering hard surface roads.

^d = log likelihood test value; ANOVA not performed on breeder survival.

Caption 11. Florida scrub-jay nest are often concealed in myrtle oaks (*Quercus myrtifolia*) with an outer structure of twigs and an inner lining of fibers pulled off cabbage palms (*Serenoa repens*).



Dispersal. Mean natal dispersal distances were 1.0 km (n = 56) for males and 2.8 km for females (n = 65); these differences were significant (p = 0.010). The longest distance dispersals were 31 km for females and 13 km for males. Ninety-five percent of

all male and 92% of all female natal dispersal traversed only 7 territories (Figure 9). Florida Scrub-Jays from the same territory cluster filled 91% of breeding vacancies that were filled by color-banded birds. We observed movements between most subpopulations in South Brevard (Table 13). Three exchanges between Central and South Brevard metapopulations were observed but no exchanges were observed between the North Brevard metapopulation Central or South Brevard metapopulations. Scrub-Jays that immigrated into the study tracts from unknown locations. Twelve of these immigrants from unknown locations became residents at Valkaria and half of all the immigrants were one-year-olds. We were unable to colorband many Florida Scrub-Jays inhabiting several study sites so that 5-20 unbanded immigrants filled vacancies. Only one Florida scrub-jay, colorbanded in our study sites, was found in a habitat fragment outside of our study sites during our peripheral surveys. The remaining exchanges occurred within our study sites.

Figure 20. Mean Dispersal Distance Between the Natal Territory and the Territory Where a Florida Scrub-Jay First Became a Breeder.




Table 13. Exchanges of Florida Scrub-Jays Among Territory Clusters.

Cluster of Origin	Destination Cluster								
	Viera	Wickam	Stewart	Melbourne Airport	Malabar-Jordan-Valkaria	Palm Bay	West Micco	Sebastian	Fellsmere
Viera									
Wickam									
Stewart									
Melbourne Airport		1							
Malabar-Jordan-Valkaria		3					1	1	1
Palm Bay					1				1
West Micco					3			1	3
Sebastian						1			2
Fellsmere							3	1	

Notes: Entries are the number of individuals

Caption 12. Optimal primary ridge (xeric oak scrub).



4.3 Discussion

Basic Demography. By 2001, demographic studies sampled 64% of the combined metapopulations of North Brevard, Central Brevard, and South Brevard-Indian River-St. Lucie. Sample sizes for survival and demographic performance were smaller because of difficulties in capturing all birds. We estimate that approximately 85% of territories were surveyed for dispersed individuals during 2001-2002. Surveys by us, U. S. Fish and Wildlife Service personnel, Brevard County and Indian River County government employees, consultants, and interested citizens suggested that few jays from our demographic study tracts successfully became breeders outside our study tract without our detection.

We estimated a 34% decline in the number of breeding pairs between 1992 and 2002 in Section 3.0. Population models based on demographic data collected from elsewhere and applied to habitat quality associated with the population studied here predicted a population decline of 40% per decade (Breininger et al. 1996b, Root 1998, Breininger 1999). All demographic variables were below levels associated with optimal habitat needed to sustain the populations (Woolfenden and Fitzpatrick 1984, 1991; Breininger et al. 1999a). Based on habitat quality and the extent of overgrown scrub in 1992 (Swain et al. 1995, Boyle 1996), it was reasonable to assume that the population declined by 30-50% a decade for at least 1-2 decades before 1992 without even considering the effects of habitat loss. Similar declines were associated with the federal properties (Breininger et al. 1996b).

Results of pair bond fidelity were similar to other published studies (Woolfenden and Fitzpatrick 1984, Breininger et al. 1996a). The frequency of both breeders dying during the same year was higher in this study because of the low survival associated with the presumed epidemic that occurred in 1997 (Breininger et al. 1999b). Rates of breeding by one-year-olds were much greater than associated with stable populations where breeding vacancies are scarce (i.e., Woolfenden and Fitzpatrick 1984). Higher rates of delayed breeding have been reported before in declining populations and delayed breeding appears directly related to breeding opportunities (e.g., Breininger et al. 1996a, Breininger 1999).

Mean family size was lower than typical for optimal habitat (Woolfenden and Fitzpatrick 1984). Increased breeding opportunities for nonbreeders occurred because mortality exceeded reproductive success and because the amount of suitable habitat increased from habitat restoration and wildfires in many areas. Habitat restoration can increase breeding opportunities as was evident by many one-year-olds that bred at restored sites at Valkaria and Sebastian Buffer Reserve.

There was much suitable but vacant habitat and a surplus of potential breeders (helpers) indicating that nonbreeders did not distribute themselves in proportion to the breeding opportunities that existed. Although much vacant habitat was marginal, there were many vacant optimal territories at Valkaria, Micco, Babcock, and north Sebastian Buffer Reserve. Helpers might choose to not disperse into suboptimal habitats but this does not explain vacant optimal territories. There are many possible explanations about why territories that recently became optimal were unoccupied for periods of time. These optimal territories were rarely adjacent to territories with helpers and were rarely near territories with helpers. Low dispersal propensities by males and conspecific attraction were two explanations. Conspecific attraction might be important because

jays from many territories provide greater vigilance against predators. Florida Scrub-Jays have a sentinel system keenly adapted to detecting accipiters (McGowan and Woolfenden 1989), which are always common along the Atlantic coast (Breininger et al. 1996a, authors personnel observations). Another explanation for unoccupied territories was that many helpers were closely related within fragments, and Florida Scrub-Jays have tendencies to avoid incest (Woolfenden and Fitzpatrick 1984). We plan to use an information-theoretic approach to test alternative models as the data is prepared for journal publication. Practical applications of these results are that attempts should be made to prevent extinction within PRUs that already have jays and prioritize restoration near territories that have helpers.

There is little data so far on Florida Scrub-Jay population responses to restoration across a broad range of conditions. Population modeling also suggested that providing opportunities for enhancing the survival of existing Florida scrub-jays and providing breeding opportunities for helpers were important because population expansion attributed to reproductive success exceeding mortality could be a slow process (Breininger et al. 1999a). This occurs partially because experienced breeders and breeders with helpers have greater demographic success than novice breeders and breeders without helpers (also see Woolfenden and Fitzpatrick 1984, 1996). Recently colonized areas tended to be inhabited by inexperienced jays. Delays in the saturation of restored areas might occur because it takes years for experienced breeders and nonbreeders to accumulate (Breininger et al. 1999aa).

Demography and Habitat Arrangements. The results were consistent with long-term studies on KSC where primary and secondary territories functioned as population sources when habitat was optimal; tertiary territories were population sinks (Breininger and Oddy 2001). Here, population sources refer to areas that produce an excess of potential breeders and emigration that exceeds immigration. These should not be confused with habitat fragments that have mortality that exceeds reproductive success but supply immigrants to other areas, such as larger habitat fragments (Breininger 1999, Bowman unpublished). Suburban areas provide an example of the latter where dispersal has been one-way: from suburbs into natural landscapes (Thaxton and Hingtgen 1996; Bowman unpublished). This latter process can only be sustained as long as suburbs have immigrants to supply. Here, population sinks refer to areas with mortality that exceeds reproductive success that can only be sustained by immigration, which exceeds emigration.

In this study, most primary territories had too much tall scrub. Oak scrub does not burn as readily as palmetto-dominated scrub (Breininger et al. 2002). Secondary territories often have greater habitat quality than oak-dominated areas because they burned more readily. This study showed that not only does palmetto-oak provide for an overall larger population but palmetto-oak in optimal condition serves as a population source. Although much xeric oak scrub was unoccupied because it was overgrown, xeric oak can contribute most to recovery once restored because territory quality generally increases as oak cover increases if other habitat conditions are not suboptimal (e.g., Breininger et al. 1995, Grubb et al. 1998, Breininger and Oddy unpublished manuscript). Although this study was too short with too few optimal territories to compare dispersal propensities among source and destination territories,

we did observe many movements from optimal to suboptimal territories. We demonstrated such source-sink dynamics within landscapes elsewhere (Breininger and Oddy 2001, Breininger and Carter 2003).

Results suggested that territories enclosed within suburbs and along roads had mortality that exceeded reproductive success, which was consistent with other studies (Breininger 1999, Mumme et al. 1990, Bowman unpublished). However, negative demographic performance might have occurred because there were too many tall shrubs and trees. Our new findings suggested that edge territories (territories in reserves along edges of human cultural landcover types but not roads) that were optimal (primary or secondary with an optimal arrangement of scrub height) were not population sinks and even functioned as population sources. We expect there could be a broad range of variation among reserve edges because of the complexity of possible edge effects (Mumme et al. 2000, Bowman and Woolfenden 2001, Schoech and Bowman 2001, Reynolds et al. 2003, Fleishcer et al. 2003, Reynolds et al. in press, Thorington and Bowman in press). Although managers often prefer to leave buffers along edges, we emphasize the importance in managing habitat to be optimal beginning at the border of potential reserves because of the relatively low number of core territories in most large reserves.

We agree with Bowman (2001; personal communication) that our poor understanding of edge-effects is one of the most important data gaps for conservation planning. Bowman (unpublished) showed that even areas with low human population densities were population sinks that could only persist with immigration. Given that jays tend not to immigrate into many of these areas from reserves (Thaxton and Hingtgen 1996, Bowman unpublished), much of the entire range of the Florida scrub-jay could collapse (Bowman 2001). If this is widespread and cannot be changed, we are purchasing conservation lands in many of the wrong areas because most areas lack enough core territories to sustain losses along the edges of reserves. If true, we should only be performing Florida scrub-jay conservation in the largest of the remaining potential scrub reserves that have an abundance of core territories.

One reason why areas with low human housing densities were population sinks in Bowman's study was that habitat quality was poor because much habitat was tall and unburned, and it was not possible to partition variance in habitat quality effects from additive effects associated with the human landscape (Bowman unpublished). Our study had potential to partition the variances associated with different factors because many territories occurred within suburbs, because a large number of territories occur in potential reserves adjacent to different types of human edges, and because many territories occurred distant from edges. To date, only limited conclusions can be made because many reserves only approached optimal at the end of the study. We emphasize caution in drawing conclusions that are broadly applicable.

The arrangement of height within territories was clearly the most influential factor influencing demographic success. The results were consistent in suggesting that territories with only short or only tall scrub were population sinks (Woolfenden and Fitzpatrick 1984, 1991; Breininger and Oddy 2001, Breininger and Carter 2003). The results for tall mixed scrub depended on landscape context. Primary and secondary territories with tall mixed scrub had reproductive success that matched the mortality rate. This differed from studies on KSC that indicate that tall mixed scrub is a

population sink (Breininger et al. 1998a, Breininger and Carter 2003) but is similar to studies on Cape Canaveral Air Station (Stevens and Young 2002). We suspect that other habitat factors, such as openings and the abundance of xeric soils might explain these differences because of compensatory effects (Burgman et al. 2001).

Nevertheless, recovery depends on population expansion, which requires a large number of territories that produce an excess of breeders. Therefore, management should not be judged as successful when there is an abundance of tall mixed scrub.

Almost all territories occurred in suboptimal habitat, which explained the population decline. However, Florida Scrub-Jays that resided in optimal habitat were successful and a large number of territories in optimal habitat could sustain a large number of territories in suboptimal habitat making potential populations large enough to withstand catastrophic events such as epidemics (Root 1998, Breininger et al. 1999aa, Stith 1999, Breininger and Oddy 2001, Breininger and Carter 2003). Habitat arrangement clearly influenced population dynamics, but our abilities to partition variance among many variables were limited by a low sample size of optimal territories. Sample sizes of optimal territories began increasing in 2001 and 2002 within many territory clusters. Partitioning variance among many potential factors (oak cover, height, tree densities, edge effects, experience, presence of helpers, annual variation) is an important investigation but we believe such an investigation should focus on territories that approach optimal and not where many habitat variable are marginal.

All evidence indicates that recruitment only exceeds mortality in recently burned areas with enough optimal height oak scrub and low tree densities (Woolfenden and Fitzpatrick 1984, 1991; Breininger and Oddy 2001; Breininger and Carter 2003). Long-term persistence depends on population recovery to increase population size within conservation areas. Population growth requires recruitment to significantly exceed mortality and therefore requires optimal height arrangements.

Dispersal Mean dispersal distances were greater than those in unfragmented landscapes (Woolfenden and Fitzpatrick 1984; Breininger and Carter 2003), as expected (Thaxton and Hingtgen 1996, Breininger 1999, Fitzpatrick et al. 1999). In just two years of study in Central Brevard we observed exchanges with South Brevard. Based on the dispersal distances observed and the presence of potential habitat, we eventually would have expected rare exchanges between Central Brevard and North Brevard.

Dispersal in Florida Scrub-Jays might best be explained by two opposite strategies: “delay and foray” and “depart and float” (Stith 1999). The delay and foray strategy explains most dispersals where nonbreeders make short forays to investigate vacancies within an assessment sphere. The depart and float dispersal strategy is most common among birds, but not among Florida Scrub-Jays. The depart and float strategy entails greater risks, especially if unsuitable habitat must be crossed (Stith 1999). We observed 16 Florida Scrub-Jays that we described as floaters. Most of these occurred in Palm Bay and seemed to remain in Palm Bay after the 1997 epidemic. We also observed a few floaters for a few years in very tall habitat surrounding Valkaria airport. Here, we describe floaters as birds that were never seen associated with a particular territory but known to be alive because they were occasionally sighted in different locations. These individuals were rarely observed to have left the territory cluster.

Many of the birds that left a cluster to become breeders in another cluster were not observed to be floaters but disappeared for several months and were re-sighted in another cluster. Many of these left clusters that had vacant territories suggesting that some exchange among territory clusters is likely.

We observed only two emigrants from the relatively large Palm Bay population, which resided in poor quality suburbs that we thought would be a source of immigrants into reserves. We observed two movements from reserves into urban fragments. Each of these came from reserve scrub patches that had few contiguous territories. Unbanded jays that immigrated into our study tracts were usually very tame upon arrival and easily captured leading us to believe they were not any of the unbanded residents we were frequently attempting to capture. We believed that many of the immigrating jays came from suburbs because they were very tame. Jays in residing in large scrub tracts areas were usually unfamiliar with peanuts and cautious at first.

There are many ideas about factors that influence dispersal rates dispersal among populations (e.g., density-dependence [Akçakaya 2002], metapopulation theories [Harrison and Taylor 1987, Hanski 1999] and ideas that it is adaptive for small populations to have greater exchange of individuals than large populations [McPeck and Holt 1992, Doncaster et al. 1997, Diffendorfer 1999]. We have not yet tested these alternatives because of many possible alternative models and the lack of a large sample size of exchanges.

We emphasized demographic responses to fragmentation because the genetic consequences are usually less immediate threats (Holsinger and Vitt 1997). There is a large gap in understanding differences between the movement of individuals and genes (Antonovics et al. 1997; Sork et al. 1998). Across long time periods, each metapopulation on the mainland of Florida's Atlantic Coast might be too small if their populations continue to decline and extirpation occurs within the intervening fragments. Theoretical models suggest that a small amount of population exchange can substantially reduce these threats (Efford 1996). Our results suggested a small amount of exchange occurs. Even a small amount of exchange influenced the colonization of territories following restoration. Based on the exchange observed and the slow pace of acquiring and restoring reserves to optimal habitat quality, we do not recommend translocation. The exception, however, occurs north of Buck Lake where there is an abundance of potential territories and very few Florida Scrub-Jays available for recolonization. Translocation there might be an important option after habitat is acquired and restored.

5.0 South Beaches Metapopulation Study Update


The purpose of this section is to update relevant material and not summarize what is examined in detail elsewhere (Breininger 1999) and used to support recovery team population modeling (Stith 1999). Current population size includes only 2 breeding pairs (Figure 11) suggesting that extinction is inevitable. A pertinent finding of our South Beaches study was that we observed frequent dispersals among population clusters and subpopulations and at least one dozen exchanges where Florida Scrub-Jays must have crossed > 4.2 km of contiguous urban areas. An interesting observation was that Florida Scrub-Jays in the Coconut Point EELs reserve had been extirpated but was

recolonized shortly after it was burned. The male that recolonized the site was from another subpopulation indicating that recolonization by males can occur. Extirpation probably again occurred because the habitat quality was poor, a busy road intersected the reserve, and the overall population size was very small.

6.0 Management Recommendations

Habitat management Fires will rarely produce a landscape with only optimal territories but recovering populations requires maximizing the proportion of optimal territories. Habitat management and restoration must address habitat suitability at the territory scale because territories are the functional demographic units within landscapes (Breininger and Carter 2003). Most populations declined because there were too many marginal territories that included tall scrub. Because tall patches are difficult to eliminate by mosaic fires without first cutting them, cutting must be applied to specific patches of tall scrub throughout most of the landscape or an extensive fire must be applied that burns nearly all scrub. Florida Scrub-Jay population declines can occur after extensive fires but these can be followed by population growth (Breininger and Oddy 2001). Extensive fires occurring once every 20 years might be useful because they burn nearly all areas, including those resistant to fire (Breininger et al. 2002). Results here suggested that short territories might have greater negative demographic success than tall mix territories and tall mix territories might have recruitment that matches mortality. The demographic success in tall mix territories appears to be site specific. If so, extensive burns in areas where the population is near carrying capacity might be disadvantageous if tall mix territories are not sinks and have much greater demographic success than short territories. Extensive burns might be essential in areas where tall mix territories are sinks (Breininger and Carter 2003). Population sizes are usually well below carrying capacity so that initial aggressive approaches are recommended to expedite restoration to a landscape of optimal territories as quickly as possible. Florida Scrub-Jays can defend relatively large areas so that extensive burns may have little negative impact in areas with few territories relative to habitat potential but the long-term benefit of extensive fires might be substantial as the population recovers.

Figure 21. Location of Florida Scrub-Jay Territory Clusters, North (N) and South (S) Subpopulations on the Barrier Island of South Brevard County (Breininger 1999). The Numbers in Parentheses Refer to the Number of Breeding Pairs from 1992 to 2002.



Once restored, frequent mosaic fires are the best mechanism to maintain openings and still keep a few acres of oak scrub at 120-170 cm within each territory (Breininger and Oddy 2001, Breininger et al. 2002). Open sandy areas are microhabitats important to Florida Scrub-Jays (Woolfenden and Fitzpatrick 1984, Breininger et al. 1995) and many specialized plants (Menges and Hawkes 1998). Most open sandy areas do not persist in scrubby flatwoods for more than a few years after fire (Schmalzer and Hinkle 1992a, Hawkes and Menges 1996). Frequent fires that originate in mesic flatwoods or marshes and burn into oak scrub represent one mechanism to retain open sandy areas. Conventional wisdom holds that natural fires frequently occur within Florida landscapes but that fires often burn out upon entering scrub oaks where flammability is lower (Webber 1935, Myers 1990, Breininger et al. 2002).

Many landscapes have become forests or otherwise have too many tall trees so that timbering also is an option. Reserve edges should also be optimally managed.

Restoring and managing scrub for Scrub-Jays will benefit other species of conservation concern and is not single species management (Means and Campbell 1981, Speake et al. 1978, Auffenberg and Franz 1982, Layne 1990, Breininger and Smith 1992, Menges and Kohfeldt 1995, Hawkes and Menges 1996, Menges and Hawkes 1998).

Reserve Design Conservation reserve designs should include secondary and tertiary ridges and adjacent mesic flatwoods as potential habitat, because they contribute to population stability by making overall populations larger (Pulliam 1988, Howe et al. 1991). Secondary and tertiary ridges are often not specified as part of the scrub ecosystem (Schmalzer et al. 1999) and are often excluded from Florida Scrub-Jay conservation reserve designs (Root 1998). Mean demographic rates provided herein can approximate the proportion of sink territories that can be supported by source territories, although these approximations are probably best done using population models that consider additional complexities, such as stage-specific vital rates, environmental stochasticity, and catastrophes (Burgman et al. 1993).

It is not necessary to explicitly map scrub oak ridges to apply these results. One can grid landscapes into potential territories using geographical information systems and categorize the habitat potential of individual grid cells to determine the potential arrangements of sources and sinks in conservation reserve designs.

Below we provide a general comment on the relative importance of every PRU. Although many sites by themselves might not be designated high priorities, recovery requires conservation of most of the units to achieve population viability. Priorities (Figure 22) assumed that it was most important to maximize the number of contiguous territories and connectivity among reserves while keeping the proportion of fragmented territories to a minimum. The draft recovery plan provides further guidance.

PRU 1. Much was forested prior to the 1998 wildfires, which burned large areas. Although most of the unit is mesic, there is potential for 10 core territories. Surveys might have underestimated population size because of its and limited access although we doubt that many families occur because it was forested before 1998. Habitat potential and possibly Florida Scrub-Jay population size have been underestimated in

South Volusia County. The possibility of connecting the North Brevard metapopulation to the Merritt Island-S.E. Volusia metapopulation should be explored.

PRU 2. Much of this scrub borders a highway and it can only support a couple pairs. Much of the site is oak scrub but its small size and lack of core territories does not make it a priority for conservation.

PRU 3. Much was forested before the 1998 wildfires, which burned large areas. The unit has a potential population size of 30 territories, including a large number of core territories. Surveys might have underestimated population size because of limited access. We doubt that many families occur there because it was forested before 1998. Acquisition of this site should be of high priority to maintain it as a large natural landscape. It occurs within a region of high conservation potential determined by Florida Greenways studies (Tom Hctor, University of Florida, personal communication). This might be a good site, combined with Buck Lake, for a translocation experiment.

PRU 4. This site is east of PRU 3 and is mostly mesic but could provide habitat for several core territories. Population size in 1992 might also have been underestimated. We recommend that the conservation potential of this site be closely evaluated, because it is not in an existing conservation proposal. We recommend the site for Florida Scrub-Jay conservation as a moderate priority. It will be important to acquire most or all of the site to maximize its potential value given that it is separated from PRU3 by a highway and bounded to the east by suburbs. .

PRUs 5, 6. This area has potential to support a few territories in a fragmented landscape.


PRU 7. This area has potential to support only 1-2 territories in a landscape fragmented by agriculture.

PRU 8. This area has potential to support several territories in a fragmented landscape but it has potential connective value.

PRU 9. This area includes the Buck Lake Conservation Area that burned in 1998 and has been subject to many restoration efforts since then. Some habitat remains available for purchase within this area. We consider this additional purchase to have high priority because of the importance of sustaining core populations and because the population sizes in northern Brevard are precariously close to extinction. This would be a good site for translocation because population size has dropped to 2 territories and there are few potential colonists nearby.

PRU8. This includes scrub in the southern end of Buck Lake Conservation Area and is badly overgrown but could be an important link to Seminole Ranch Conservation Area. Some additional habitat could be purchased but it would be almost surrounded by human development. Restoration of scrub in conservation ownership should be a top priority.

Figure 22. Possible Reserve Design.



PRUs 11, 12, 13. These units have territories within a fragmented landscape.

PRU 14. Most of this unit is part of Seminole Ranch Conservation Area badly needs habitat management.

PRU 15. This unit could support 17 territories. Most of this unit is part of Seminole Ranch Conservation Area but it badly needs habitat management, which should be a top priority. The Florida Fish and Wildlife Conservation Commission is the lead agency responsible for habitat management. Purchasing the remaining areas should be a top priority because many would be core territories and they would maximize potential population size, which is needed to sustain population viability.

PRUs 16, 17. These habitat fragments, surrounded by suburbs, can only support a few territories.

PRU 18. The area South of South Lake and mostly west of Fox Lake is within a proposed CARL boundary and could support about 20 territories in a predominately natural landscape connected to the Seminole Ranch Conservation area. Therefore, this should be a top priority for conservation.

PRU 19. This small habitat fragment can support 5 territories.

PRU 20. This area includes the Titusville Well-Field and a site purchased by EELs. There are some areas that still should be acquired. Most of the site has not been managed. Restoration should be a top priority given that it could support about 14 territories but only 2 remain occupied by Florida Scrub-Jays.

PRU 21. Some of this area is within proposed and existing EELs sanctuaries, including the Enchanted Forest. It has potential to support 17 territories and has connective value, although suburbs or forest would border most territories. We suggest it is of moderately high priority even though most of it is unoccupied.

PRU 22. This fragment is surrounded by suburbs and highways and has potential to support contiguous 14 territories but only two occupy the site.

PRU 23. This area is surrounded by suburbs but could support one territory.

PRU 24. The area surrounding Tico Airport has potential to support 55 territories. Acquiring large portions of this area should be a top priority because of the large potential population size and its connective value.

PRUs 25, 26, 27, 28, 30. These habitat fragments are bordered by roads but have potential habitat to support several territories.

PRU 29. This area along Grissom Road provides habitat for about 30 territories, although road mortality along roads is a concern. A portion of this area has been bought for conservation under the Grissom Road Mega-Parcel.

PRU 31. This area along Grissom Road and north of Cocoa is the southern terminus of the North Brevard population making it a top priority to maintain opportunities for exchange with the larger southern metapopulations. It has potential to almost 20 territories but has not been proposed for conservation.

PRU 32. This area surrounding Mud Lake has potential to support 6 territories but these potential territories are unoccupied because the area has become forest. This site has local conservation interest because it is one of the few natural areas remaining in the City of Cocoa. It could have value to connect metapopulations.

PRUs 33, 34, 35. These small fragments could support several territories but none are known to be occupied, although these could be restored and might have value in linking metapopulations.

PRU 36. This is the single most critical polygon in Central Brevard because it is inhabited by one the largest populations and it has some of the best habitat quality. Almost half is part of the EELs Cruickshank Sanctuary and the Viera Mitigation Site. Significant portions are at great risk to development. Given this is the critical core for Central Brevard, the opportunity to link metapopulations probably depends upon the viability of this unit and therefore we recommend it be given a top priority for conservation.

PRUs 37- 46, 48. The habitat fragments near Wickam Road once could support 20 territories but habitat is rapidly being lost to development.

PRU 49. This habitat fragment could support 8 territories and much or all of the habitat is part of Wickham Park. Developing a conservation plan for Wickam Park should be a top priority because of its importance in connecting metapopulations.

PRUs 49, 50, 51. These habitat fragments could support a few territories.

PRUs 52, 53, 54, 55, 56, 57. The potential population size around Melbourne Airport exceeds 20 territories making conservation of at least some of the PRUs a top high priority for linking metapopulations.

PRUs 58, 59, 60. These habitat fragments can support few territories.

PRUs 61, 62, 63. Each of these can support many territories and they are the deciding links determining whether the small and vulnerable Central Brevard is a separate metapopulation from the South Brevard-Indian River-St Lucie metapopulation based on criteria described by Stith et al. 1996. No conservation proposals address conservation in these areas but their fate needs immediate evaluation, depending on recovery plan criteria.

PRU 64. This includes the Turkey Creek and Malabar Scrub Sanctuaries. Most of the unit has been acquired; extensive habitat restoration should be a top priority.

PRU 65. This includes the Jordan scrub sanctuary and Waterside Downs mitigation area. There are about 16 additional potential unprotected territories that are a top priority because they would provide a critical link between the Malabar Scrub Sanctuary and Valkaria. The Malabar-Grant-Valkaria has the greatest potential to be the largest, contiguous remaining subpopulation along the Atlantic Coastal Ridge, which once stretched along nearly all of Florida's east coast.

PRU 66. This PRU near Brevard County Community College has potential to support a Florida scrub-jay territory.

PRU 67. This area established for scrub-jay conservation at Liberty Park could support one territory.

PRU 69. This habitat fragment east of the Palm Bay suburbs has enough habitat to support 6 territories. The site is proposed for conservation and might be an important connector. Some of it is an EELs sanctuary that badly needs management and has a private landowner interested in conservation.

PRU 68. Much habitat is public ownership although considerable amounts remain to be acquired. Conservation plans that restore more habitat at Valkaria Airport and the Habitat Golf Course are needed. Core territory integrity is uncertain because of a potential Florida Inland Navigation District (FIND) Spoil Site. This site has the

necessary permits to destroy the potential core of the Valkaria Scrub Reserve. FIND has expressed willingness to shift the site. Several of the problems in moving the site have been associated with nearby wetlands, which are disturbed and have no regional significance in contrast to the importance of maintaining this last remnant of an Atlantic Coastal Ridge Scrub Ecosystem.

PRUs 71, 72, 74-77. These habitat fragments could support a few territories.

PRU 70. The proposed Babcock Sanctuary can support 8 territories. Much of the site is proposed for conservation and has been used for mitigation. Much of the site used for mitigation is approaching optimal conditions.

PRU 73. This area can support almost 22 core territories. Much habitat south of Grant Road has been used for mitigation and has been restored. A large tract of contiguous scrub occurs just north of the Micco Scrub Sanctuary should be a top priority for conservation.

PRU 78. This might support a few territories with restoration and might facilitate exchange between Atlantic Coastal Ridge and Ten Mile Ridge populations once restored. Most has been acquired by EELs.

PRU 79. This area can >35 territories and comprises the Micco Scrub Sanctuary and North Sebastian Buffer Reserve. Restoration has been completed on Sebastian Buffer but additional mechanical cutting and burning is still needed on the Micco Scrub Sanctuary.

PRU 80. The area south of Micco could support 10 territories and has moderately important value in connecting subpopulations.

PRUs 81, 82. These areas on north Sebastian Buffer Reserve could several territories. Restoration would need to include the intervening pine flatwoods. Restoration should have long-term value by connecting subpopulations although there is no immediate need.

PRU 83, 84. These areas on north Sebastian Buffer Reserve can support almost 20 territories but the landscape share habitat with one of the last Red-Cockaded Woodpecker colonies within the region and it is precariously close to extinction. Although habitat requirements overlap optimal pine densities differ between species. We recommend actions that only benefit both species rather than maximizing carrying capacity for Florida Scrub-Jays only. Reduction of hardwood trees near the North Fork of the Sebastian River could benefit both species.

PRU 85. The Coracii portion of Sebastian Buffer, heavily degraded by agriculture, can support >35 territories and much restoration has begun. Most territories remain marginal because the recently burned scrub is interspersed by forest and tall scrub. The Sebastian Area-Wide Florida Scrub-Jay Habitat Conservation Plan (Sebastian HCP) reserves occur within dispersal distance of Coracii. A wide open landscape should be restored along much of the northern and eastern border to enhance exchange with the northern part of the buffer and the Sebastian Highlands.

PRU 86. Detailed state-of-the-art management plans were developed as part of the Sebastian HCP (Smith Environmental Services 1999). This area also includes some mitigation land and areas that have not been proposed for conservation. This areas could connect PRUs 85 and 86. The entire area could support >10 territories and maximizing population size should be a top priority.

PRU 87. The Carson Platt portion of the Sebastian Buffer Reserve is the largest contiguous stretch of habitat approaching optimal habitat quality found anywhere along Florida's Atlantic coast. There is enough habitat for approximately 40 territories. Many patches of tall scrub need fire and the area would benefit most by targeting these patches and burning the rest with mosaic fire. The mesic flatwoods east and west of the scrub ridge need hot fire. There is much hardwood invasion of swales and mesic flatwoods along the northern edge that could be reduced to enhance connectivity.

PRU 88. A local park and some additional overgrown scrub along the South Fork could be restored to enhance connectivity between Sebastian Airport and Sebastian Buffer Reserve.

PRU 89, 90, 91, 93-96. These PRUs include the Sebastian Highlands HCP. Several additional properties (Water Management District, Wabasso Golf Course) that could be acquired or managed to enhance the HCP. Populations in these areas precariously close to extinction.

PRU 92, PRU 97-101. A top priority should be given to acquire and restore all of these units as a contiguous scrub and flatwoods landscape separated by Interstate 95. The oak and palmetto-oak is not contiguous so that the landscape was subdivided into units for prioritizing each of them. The number of territories that could be supported is approximately 50. The only areas in conservation include the Sand Lakes Restoration Area, which can support 5 territories. The SJRWMD has been restoring this landscape (Sean Rowe personnel communication). No other conservation lands proposals include these areas so that developing acquisitions should be a top priority. The most important units are PRU 92 and 98-100.

Acknowledgements

We thank Dawn Zattau of the U. S. Fish and Wildlife Service for all her encouragement. Keith Fisher and the Florida Department of Environmental Protection provided additional support. Dale Wade of the Southern Forest Research Station also provided additional support through the Florida Fire Science Team. We thank Anne Birch of the Brevard County Environmentally Endangered Lands Committee, Peter Henn of the St. Johns River Water Management District, and Zeke Nations of Endangered Lands Management Corporation for assisting us with access issues. Brad Stith, Reed Bowman, Glen Woolfenden, Paul Schmalzer, and Brean Duncan provided an abundance of helpful advice.

Literature Cited

- Akçakaya, H.R. 2002. RAMAS GIS: linking landscape data with population viability analysis (version 4.0). Applied Biomathematics, Setauket, New York.
- Akçakaya, H. R., Burgman, M. A., and Ginzburg, L. R. 1999. Applied Population Ecology. Second Edition. Sunderland, Massachusetts: Sinauer Associates.
- Antonovics, J., Thrall, P. H., and Jarosz, A. M. 1997. Genetics and the spatial ecology of species interactions: the *Silene-Ustilago* system. In *Spatial Ecology: The Role of Space in Population Dynamics and Interspecific Interactions*, eds. D. Tilman and P. Kareiva, pp. 158-184. Princeton: Princeton University Press.

- Auffenberg, W. and R. Franz. 1982. The status and distribution of the gopher tortoise (*Gopherus polyphemus*). Pages 95-126 in R.B. Bury, editor. North American tortoises: conservation and ecology. U.S. Fish and Wildlife Service Research Report 12.
- Boyle, S. R. 1996. A landscape approach to Florida scrub-jay (*Aphelocoma coerulescens*) habitat use in Brevard County, Florida. M. S. Thesis, Florida Institute of Technology, Melbourne, Florida. 134 pp.
- Bowman, R. and G. E. Woolfenden. 2001. Nest success and the timing of nest failure of Florida Scrub-Jays in suburban and wildland habitats. Pages 383-402, In Avian Conservation and Ecology in an Urbanizing World, Marzluff, J.M., R. Bowman, and R.E. Donnelly (eds.). Kluwer Academic Publ., New York, NY.
- Breiner, D. R. 1999. Florida Scrub-Jay demography and dispersal in a fragmented landscape. *Auk* 116:520-527.
- Breiner, D. R. and G. M. Carter. 2003. Territory quality transitions and source-sink dynamics in a Florida Scrub-Jay population. *Ecological Applications* 13:516-529.
- Breiner, D. R. and D. M. Oddy. 2001. Fire and Florida scrub-jay source-sink dynamics in mesic flatwoods. Pp. 3-7 in D. P. Zatta, ed., Proceedings of the Florida Scrub Symposium 2001. U.S. Fish and Wildlife Service, Jacksonville, Florida.
- Breiner, D. R., and P. A. Schmalzer. 1990. Effects of fire and disturbance on plants and animals in a Florida oak/palmetto scrub. *American Midland Naturalist* 123:64-74.
- Breiner, D. R., and R. B. Smith. 1992. Relationships between fire and bird density in coastal scrub and slash pine flatwoods in Florida. *American Midland Naturalist* 127:223-240.
- Breiner, D. R., B. W. Duncan, and N. J. Dominy. 2002. Relationships between fire frequency and vegetation type in pine flatwoods of east-central Florida, USA. *Natural Areas Journal* 22:186-193.
- Breiner, D. R., M. A. Burgman, and B. M. Stith. 1999a. Influence of habitat, catastrophes, and population size on extinction risk on Florida Scrub-Jay populations. *Wildlife Society Bulletin* 27: 810-822.
- Breiner, D. R., M. L. Legare, and R. B. Smith. In press. Edge effects and population viability of Eastern Indigo Snakes in Florida. Pages 000-000 in H. R. Akçakaya, M. Burgman, O Kindvall, P. Sjorgren-Gulve, J. Hatfield, and M. McCarthy, editors. Species conservation and management: case studies. Oxford University Press.
- Breiner, D. R., M. J. Provanca, and R. B. Smith. 1991. Mapping Florida scrub jay habitat for purposes of land-use management. *Photogrammetric Engineering and Remote Sensing* 57: 1467-1474.
- Breiner, D. R., B. A. Burgman, H. R. Akçakaya and M. O. O'Connell. 2001. Use of metapopulation models in conservation planning, K. J. Gutzwiller, editor. Pages 405-427 in *Concepts and Applications of Landscape Ecology in Biological Conservation*. Springer-Verlag, New York.
- Breiner, D. R., V. L. Larson, B. W. Duncan, R. B. Smith. 1998a. Linking habitat suitability to demographic success in Florida Scrub-Jays. *Wildlife Society Bulletin* 26:118-128.

- Breininger D. R., V.L. Larson, D. M. Oddy, R. B. Smith. In press. How does variation in fire history influence Florida scrub-jay demographic success? In press: Fire effects on rare and endangered species conference. International Association of Wildland Fire, Fairfield, Washington.
- Breininger, D. R., D. M. Oddy, M. L. Legare, and B. W. Duncan 1999b. Developing biological criteria for the recovery of Florida scrub-jay populations on public lands in Brevard County: patterns of fire history, habitat fragmentation, habitat use and demography. In J. D. Brenner (editor). Ecological and economic consequences of the 1998 Florida wildfires: http://flame.fl-dof.com/joint_fire_sciences/.
- Breininger, D. R., V. L. Larson, D. M. Oddy, R. B. Smith and M. J. Barkaszi. 1996a. Florida scrub-jay demography in different landscapes. *Auk*: 112:617-625.
- Breininger, D. R., V. L. Larson, B. A. Duncan, R. B. Smith, D. M. Oddy, and M. Goodchild. 1995. Landscape patterns in Florida Scrub Jay habitat preference and demography. *Conservation Biology* 9: 1442-1453.
- Breininger, D. R., V. L. Larson, R. Schaub, P. A. Schmalzer, B. W. Duncan, D. M. Oddy, R. B. Smith, F. Adrian and H. Hill, Jr. 1996b. A Conservation Strategy for the Florida scrub-jay on John F. Kennedy Space Center/Merritt Island National Wildlife Refuge: an initial scientific basis for recovery. NASA Technical Memorandum 111676.
- Burgman, M. A. S. Ferson, & H. R. Akcakaya. 1993. Risk assessment in conservation biology. Chapman and Hall, London.
- Burgman, M. A., D. R. Breininger, B. W. Duncan, and S. Ferson. 2001. Setting reliability bounds on habitat suitability indices. *Ecological Applications* 10: 70-78.
- Dias, P. C. 1996. Sources and sinks in population biology. *Trends in Evolutionary Ecology* 11:326-330.
- Diffendorfer, J. E. 1998. Testing models of source-sink dynamics and balanced dispersal. *Oikos* 81:417-433.
- Doak, D. F., and Mills, L. S. 1994. A useful role for theory in conservation. *Ecology* 75:615-626.
- Doncaster, C. P., Clobert, J., Doligez, Gustafsson, L., and Danchin, E. 1997. Balanced dispersal between spatial varying local populations: an alternative to the source-sink model. *Amer. Natur.*150:425-445.
- Drechsler, M. and Wissel, C. 1998. Trade-offs between local and regional scale management of metapopulations. *Biol. Conserv.* 83:31-41.
- Duncan, B.W. and P.A. Schmalzer. In press. Anthropogenic influences on potential fire spread in a pyrogenic ecosystem of Florida. *Landscape Ecology*.
- Duncan, B. W., D. R. Breininger, P. A. Schmalzer, and V. L. Larson. 1995. Validating a Florida Scrub Jay habitat suitability model, using demography data on Kennedy Space Center. *Photogrammetric Engineering and Remote Sensing* 56:1361-1370.
- Duncan, B. A., S. Boyle, D. R. Breininger, and P. A. Schmalzer. 1999. Landscape Ecology. Coupling past management practice and historical landscape change on John F. Kennedy Space Center. *Landscape Ecology* 14:291-309.
- Dye, R. 1991. Use of firing techniques to achieve naturalness in Florida Parks. *Proceedings of the Tall Timbers Fire Ecology Conference* 17:353-360.

- Efford, M. G. 1996. Simulating spatially distributed populations. Pages 409-418 in R. B. Floyd, A. W. Sheppard, and P. J. De Barro, editors, *Frontiers of Population Ecology*. Canberra: The Commonwealth Scientific and Industrial Research Organization.
- Fitzpatrick, J. and G. E. Woolfenden. 1986. Demographic routes to cooperative breeding some New World jays. Pages 137- 160 in Nitecki and J. J. Kitchell, editors. *Evolution of behavior*. University of Chicago Press, Chicago, Illinois, USA.
- Fitzpatrick, J. W., G. E. Woolfenden, and M. T. Kopeny. 1991. Ecology and development-related habitat requirements for the Florida scrub jay (*Aphelocoma c. coerulescens*). Florida Game and Fresh Water Fish Commission Nongame Wildlife Report No. 8. Tallahassee, Florida. 49pp.
- Fitzpatrick, J. W., G. E. Woolfenden, and R. Bowman. 1999. Dispersal distance and its demographic consequences in the Florida scrub-jay. Pages 2465-2479 in Adams, N. J. and R. H. Slotow, editors. *Proceedings of the 22 International Ornithological Conference*. Durban, South Africa.
- Fleischer, A. L., R. Bowman, G. E. Woolfenden. In Press. Variation in foraging behavior, diet, and time of breeding of Florida scrub-jays in suburban and wildland habitats. *Condor*.
- Garshelis, D. L. 2000. Delusions in habitat evaluation: measuring use, selection, and importance. Pages 111-164 in L. Boitani and T. K. Fuller, editors. *Research techniques in animal ecology*, Columbia University Press, New York, USA.
- Greene, C. M. and J. A. Stamps. 2001. Habitat selection at low population densities. *Ecology* 82:2091-2000.
- Grubb, T. C. Jr., G. E. Woolfenden, and J. W. Fitzpatrick. 1998. Factors affecting nutritional condition of fledgling Florida Scrub-Jays: a ptilochronology approach. *Condor* 100:753-756.
- Hanski, I. 1999. *Metapopulation Ecology*. New York: Oxford University Press.
- Harrison, S., and Taylor, A. D. 1997. Empirical evidence for metapopulation dynamics. In *Metapopulation Biology: Ecology, Genetics, and Evolution*, eds. I. A. Hanski and M. E. Gilpin, pp. 27-42. San Diego: Academic Press.
- Hawkes, C. V. and E. S. Menges. 1996. The relationship between open space and fire for species in a xeric Florida shrubland. *Bulletin of the Torrey Botanical Club* 123:81-92.
- Holsinger, K. E., and Vitt, P. 1997. The future of conservation biology: what is a geneticist to do? In *The Ecological Basis for Conservation: Heterogeneity, Ecosystems, and Biodiversity*, eds. S.T.A. Pickett, R. S. Ostfeld, M. Shachak, and G. E. Likens, pp. 202-216. New York: Chapman and Hall.
- Howe, R. W., G. J. Davis, and V. Mosca. 1991. The demographic significance of "sink" populations. *Biological Conservation* 57:239-255.
- Huckle, H. F., H. D. Dollar, and R. F. Pendleton. 1974. Soil survey of Brevard County, Florida. U.S. Department of Agriculture, Soil Conservation Service, Washington, D. C., USA.
- Kareiva, P. 1990. Population dynamics in spatially complex environments: theory and data. *Philosophical Transactions of the Royal Society of London*. 330:175-190.

- Koenig, W. D., Philip N. Hodge, M. T. Stanback, and J. Haydock. 2000. Natal dispersal in cooperatively breeding acorn woodpecker. *Condor* 102:492-502.
- Layne, J. N. 1990. The Florida mouse. Pages 1-21 in C.K. Dodd, Jr., R. E. Ashton, Jr., R. Franz, and E. Wester, editors. Burrow associates of the gopher tortoise. Proceedings of the 8th annual meeting of the Gopher Tortoise council. Florida Museum of Natural History, Gainesville.
- Lidicker, W. Z. 1975. The role of dispersal in the demography of small mammals. Pages 103-128 in F. B. Golley, K. Petrusewicz, and L. Ryszkowski, editors. Small mammals: their productivity and population dynamics. Cambridge University Press, Cambridge, UK.
- McGowan, K. J., and G. E. Woolfenden. 1989. A sentinel system in the Florida scrub jay. *Animal Behavior* 37:1000-1006.
- McPeck, M. A., and R. D. Holt. 1992. The evolution of dispersal in spatially and temporally varying environments. *American Naturalist* 140:1010-1027.
- Means, D. B. and H. W. Campbell. 1981. Effects of prescribed burning on amphibians and reptiles. Pages 89-90 in G. W. Wood, editor. Prescribed fire and wildlife in southern forests. Belle W. Baruch Forest Science Institute, Georgetown, South Carolina.
- Menges E,S, and C.V. Hawkes. 1998. Interactive effects of fire and microhabitat on plants of Florida scrub. *Ecological Applications* 8:935-946.
- Menges, E. S. and N. Kohfeldt. 1995. Life history strategies of Florida scrub plants in relation to fire. *Bulletin of the Torrey Botanical Club* 122:282-297.
- Myers, R. L. 1990. Scrub and high pine. Pages 150-193 in R. L. Myers and J. J. Ewell, editors. *Ecosystems of Florida*. University of Central Florida Press, Orlando, Florida.
- Mumme, R. L. S. J. Schoech, G. E. Woolfenden, and J. W. Fitzpatrick. 2000. Life and death in the fast lane: demographic consequences of road mortality in the Florida Scrub Jay? *Conservation Biology* 14:501-512.
- Nagelkerke, K., J. Verboom, F. v. D. Bosch, and K. V. D. Wolfshaar. 2002. Time lags in metapopulation responses to landscape change. Pages 330-354 in K. J. Gutzwiller, editor. *Applying landscape ecology in biological conservation*. Springer, New York, USA.
- Noss, R.F., M.A. O'Connell, and D.D. Murphy. 1997. *The Science of Conservation Planning: Habitat Conservation under the Endangered Species Act*. Island Press, Washington, D.C.
- Pulliam, H. R., 1988. Sources, sinks, and population regulation. *American Naturalist* 132:652-661.
- Pulliam, H. R. 1996. Sources and sinks: empirical evidence and population consequences. Pages 45-70 In O. L. E. Rhodes, Jr., R. H. Chesser, and M. H. Smith. *Population dynamics in ecological space and time*. University of Chicago Press, Chicago, Illinois, USA.
- Pulliam, H. R., and B. J. Danielson. 1991. Sources, sinks, and habitat selection: a landscape perspective on population dynamics. *American Naturalist* 137:S50-S66.
- Pulliam, H. R., J. B. Dunning, and J. Liu. 1992. Population dynamics in complex landscapes: a case study. *Ecological Applications* 2:165-177.

- Reynolds, S. J., S. J. Schoech, R. Bowman. 2003. Nutritional quality of prebreeding diet influences breeding performance of the Florida Scrub-Jay. *Oecologia* 2003:308-316.
- Reynolds, S. J., S. J. Schoech, R. Bowman. In press. Diet quality during prelaying and nestling periods influence growth and survival of Florida scrub-jay *Aphelocoma coerulescens* chicks. *Journal of Zoology*.
- Rodenhouse, N. L., T. W. Sherry, and R. T. Holmes. 1997. Site-dependent regulation of population size: a new synthesis. *Ecology* 78:2025-2042.
- Root, K. V. 1998. The effects of habitat quality, connectivity, and catastrophes on a threatened species. *Ecological Applications* 8:854-865.
- Schmalzer, P. A. and S. R. Boyle. 1998. Restoring long-unburned oak-saw palmetto scrub requires mechanical cutting and prescribed burning. *Restoration and Management Notes* 16:96-97.
- Schmalzer, P. A., and C. R. Hinkle. 1992a. Recovery of oak-saw palmetto after fire. *Castanea* 57:158-173.
- Schmalzer, P. A., and C. R. Hinkle. 1992b. Species composition and structure of oak-saw palmetto scrub vegetation. *Castanea* 57:220-251.
- Schmalzer, P. A., S. R. Boyle, and H. M. Swain. 1999. Scrub ecosystems of Brevard County, Florida: a regional characterization. *Florida Scientist* 62:13-47.
- Schmalzer, P. A., C. R. Hinkle, and J. L. Mailander. 1991. Changes in community composition and biomass in *Juncus roemerianus* Scheele and *Spartina bakeri* Merr. marshes one year after a fire. *Wetlands* 11:67-86.
- Schmalzer, P. A., D. R. Breininger, F. Adrian, R. Schaub, and B. W. Duncan. 1994. Development and implementation of a scrub habitat compensation plan for Kennedy Space Center. NASA Technical Memorandum 109202.
- Schoech, S. J. and R. Bowman. 2001. Variation in the timing and breeding between suburban and wildland Florida scrub-jays: do physiological measures reflect different environments. Pages 289-306-402 in *Avian Conservation and Ecology in an Urbanizing World*, Marzluff, J.M., R. Bowman, and R.E. Donnelly (eds.). Kluwer Academic Publishers, New York, NY.
- Smith Environmental Services. 1999. Sebastian area-wide Florida scrub-jay habitat conservation plan. Indian River County, Florida.
- Sork, V. L., D. Campbell, R. Dyer, J. Fernandez, J. Nason, R. Petit, P. Smouse, and E. Steinberg. 1998. Proceedings from a Workshop on Gene Flow in Fragmented, Managed, and Continuous Populations. Research Paper No. 3. Santa Barbara: National Center for Ecological Analysis and Synthesis.
- Speake, D. W., J. A. McGlincy, and T. A. Colvin. 1978. Ecology and management of the eastern indigo snake in Georgia: a progress report. Pages 64-73 in R. R. Odum and L. Landers, editors. Proceedings of Rare and Endangered Wildlife Symposium, Georgia Department Natural Resources Game and Fish Division Technical Bulletin WL4.
- SPSS, Inc. 1999. SPSS/PC+Statistics. Version 9.0. SPSS, Chicago, Illinois, USA.
- Stevens, T., and J. Hardesty. 2000. Status and distribution of the Florida scrub-jay (*Aphelocoma coerulescens*) at Cape Canaveral Air Station, Florida. 45th CES/CEV, Patrick Air Force Base, FL.

- Stevens T. and J. Young. 2002. Status and distribution of the Florida Scrub-Jay (*Aphelocoma coerulescens*) at Cape Canaveral Air Force Station, Florida. 45th CES/CEV, Patrick Air Force Base, FL.
- Stith, B. M., J. W. Fitzpatrick, G. E. Woolfenden, and B. Pranty. 1996. Classification and conservation of metapopulations: a case study of the Florida scrub jay. Pages 187-216 In D.R. McCullough (ed.) *Metapopulations and wildlife Conservation*. Island Press, Ca.
- Stith, B. M. 1999. *Metapopulation dynamics and landscape ecology of the Florida scrub-jay (Aphelocoma coerulescens)*. University of Florida, Gainesville, Florida.
- Swain, H., P. A. Schmalzer, D. R. Breininger, K. Root, S. Bergen, S. Boyle, S. MacCaffree, S. Heflick, and T. Gregory. 1995. Consultants report to the Brevard County Habitat Conservation Plan. Brevard County, Florida.
- Toland, B. 2000. Current Status and Conservation Recommendations For the Florida Scrub-Jay in Brevard County. Report to the Brevard County Board of County Commissioners, Brevard County Natural Resources Office, Viera, Florida.
- Thaxton, J. E. and T. M. Hingtgen. 1996. Effects of suburbanization and habitat fragmentation on Florida scrub-jay dispersal. *Florida Field Naturalist* 24:25-37.
- Thorington, K. K. and R. Bowman. 2003. Predation rate on artificial nests increases with human housing density in suburban habitats. *Ecography* 26:188-196.
- Van Horne, B. 1983. Density as a misleading indicator of habitat quality. *J. Wildl. Manage.* 47:813-901.
- Webber, H. J. 1935. The Florida scrub, a fire-fighting association. *American Journal of Botany* 22:344-361.
- Wettstein, C. A., C. V. Noble, and J. D. Slabaugh. 1984. Soil survey of Indian River County, Florida. USDA Soil Conservation Service.
- Woolfenden, G. E. 1974. Nesting and survival in a population of Florida Scrub Jays. *Living Bird* 12:25-49.
- Woolfenden, G. E., and J. W. Fitzpatrick. 1984. *The Florida Scrub Jay: demography of a cooperative-breeding bird*. Princeton Univ. Press, Princeton, New Jersey.
- Woolfenden, G. E., and J. W. Fitzpatrick . 1991. Florida Scrub Jay ecology and conservation. Pages 542-565 in *Bird population studies* (C. M. Perrins, J. D. Lebreton, and G. J. M. Hirons, Eds.). Oxford Univ. Press, New York.
- Woolfenden, G. E., and J. W. Fitzpatrick. 1996. Florida scrub-jay (*Aphelocoma coerulescens*). Pages 1-28 in A. Poole and F. Gill, editors. *The Birds of North America*, No. 228. The Academy of Natural Sciences, Philadelphia and The American Ornithologists' Union, Washington D.C.